

Supplemental Information

SUPPLEMENTAL TABLE 4 Characteristics of CMC With at Least 4 CCC, CMC With <4 CCC, and CSHCN With at Least 4 CCC

Characteristic	CMC and 4+ CCC (<i>n</i> = 62)	CMC and <4 CCC (<i>n</i> = 15)	CSHCN and 4+ CCC (<i>n</i> = 193)	Group Comparisons, <i>P</i>			
				Overall	Column 1 Versus Column 2	Column 2 Versus Column 3	Column 1 Versus Column 3
TA from ICD codes, <i>n</i> (%)	62 (100.0)	13 (86.7)	164 (85.0)	<.001	.04	<.99	<.001
TA from screenings, <i>n</i> (%)	62 (100.0%)	15 (100.0%)	109 (56.5%)	<.001	–	.001	<.001
No. subspecialists seen in last 12 mo, median (25th, 75th percentiles)	5 (4, 6)	4 (3, 5)	2 (1, 3)	<.001	.14	<.001	<.001
NI, <i>n</i> (%)	52 (83.9)	13 (86.7)	97 (50.3)	<.001	>.99	.007	<.001

All unique patients were screened in FY2015 to quarter 1 of FY2018 (8 in FY2018). Patients screened multiple times were assigned to most complex result. CCC, TA, and NI were based on diagnosis codes from any encounter. Values presented are counts (%) for categorical variables or median (25th, 75th percentiles) for continuous variables. *P* values for comparisons among categorical variables are from Fisher's exact test. *P* values from comparisons among >2 continuous variables are from Kruskal-Wallis test and between 2 continuous variables are from Wilcoxon rank test. ICD, *International Classification of Diseases*; –, not applicable.