

## Supplemental Information

**SUPPLEMENTAL TABLE 5** Patient and Caretaker Anxiety Linear Regression Model: T1 to T2

Group	Unstandardized Coefficients		Standardized Coefficients	<i>t</i>	<i>P</i>
	$\beta$	SE	$\beta$		
All patients					
Constant	1.099	0.152	—	—	—
Sex <sup>a</sup>	−0.034	0.080	−.044	−0.429	.669
Age, y	−0.017	0.012	−.145	−1.459	.148
Psychiatric diagnosis <sup>b</sup>	0.098	0.103	.095	0.952	.344
Previous hospitalizations	−0.032	0.059	−.055	−0.534	.595
Study group <sup>c</sup>	−0.320	0.077	−.407	−4.137	<.001
Patients aged 5–10 y					
Constant	1.215	0.356	—	—	—
Sex <sup>a</sup>	−0.178	0.143	−.188	−1.247	.219
Age, y	0.000	0.040	.001	0.004	.997
Psychiatric diagnosis <sup>b</sup>	0.202	0.192	.151	1.050	.300
Previous hospitalizations	−0.032	0.142	−.034	−0.223	.824
Study group <sup>c</sup>	−0.316	0.137	−.335	−2.303	.026
Patients aged 11–16 y					
Constant	0.587	0.213	—	—	—
Sex <sup>a</sup>	0.130	0.061	.254	2.143	.038
Age, y	0.025	0.015	.189	1.634	.110
Psychiatric diagnosis <sup>b</sup>	−0.040	0.072	−.063	−0.547	.587
Previous hospitalizations	0.017	0.018	.103	0.895	.376
Study group <sup>c</sup>	−0.304	0.059	−.607	−5.147	<.001
All caretakers					
Constant	0.986	0.055	—	—	—
Sex <sup>a</sup>	0.082	0.053	.153	1.554	.125
Age, y	−0.013	0.025	−.051	−0.510	.611
Study group <sup>c</sup>	−0.224	0.042	−.528	−5.270	<.001

—, not applicable.

<sup>a</sup> Sex code: 0 = female; 1 = male.

<sup>b</sup> Psychiatric diagnosis code: 0 = no psychiatric diagnosis; 1 = psychiatric diagnosis.

<sup>c</sup> Study group code: 0 = control; 1 = intervention.

**SUPPLEMENTAL TABLE 6** Subgroup Analysis Linear Regression Model: T1 to T2

Group	Unstandardized Coefficients		Standardized Coefficients	<i>t</i>	<i>P</i>
	$\beta$	SE	$\beta$		
Hospital naïve					
Constant	1.265	0.159	—	—	—
Sex <sup>a</sup>	−0.032	0.086	−.043	−0.373	.710
Age, y	−0.019	0.013	−.170	−1.527	.131
Psychiatric diagnosis <sup>b</sup>	0.197	0.118	.182	1.670	.100
Study group <sup>c</sup>	−0.309	0.085	−.411	−3.658	.001
Hospital experienced					
Constant	1.122	0.362	—	—	—
Sex <sup>a</sup>	−0.041	0.212	−.043	−0.195	.848
Age, y	0.001	0.030	.006	0.027	.979
Psychiatric diagnosis <sup>b</sup>	−0.166	0.226	−.172	−0.735	.473
Study group <sup>c</sup>	−0.433	0.199	−.495	−2.176	.045
No psychiatric diagnosis					
Constant	1.127	0.154	—	—	—
Sex <sup>a</sup>	0.042	0.083	.056	0.511	.611
Age, y	−0.009	0.013	−.074	−0.699	.487
Previous hospitalizations	0.038	0.075	.055	0.512	.610
Study group <sup>c</sup>	−0.343	0.081	−.451	−4.251	<.001
Psychiatric diagnosis					
Constant	1.863	0.422	—	—	—
Sex <sup>a</sup>	−0.390	0.246	−.438	−1.588	.141
Age, y	−0.043	0.029	−.386	−1.474	.169
Previous hospitalizations	−0.114	0.109	−.270	−1.050	.316
Study group <sup>c</sup>	−0.321	0.233	−.363	−1.376	.196

—, not applicable.

<sup>a</sup> Sex code: 0 = female; 1 = male.<sup>b</sup> Psychiatric diagnosis code: 0 = no psychiatric diagnosis; 1 = psychiatric diagnosis.<sup>c</sup> Study group code: 0 = control; 1 = intervention.