

Supplemental Information

SUPPLEMENTAL TABLE 5 Diagnostic Codes for Included Autoimmune Diseases Based on the WHO ICD Codes From the ICD-8 and the ICD-10

Autoimmune Diseases	ICD-8	ICD-10
Alopecia areata	704.00	L63
Ankylosing spondylitis	712.49	M45
Autoimmune hemolytic anemia, acquired hemolytic anemia	283.90, 283.91	D59.1
Other hereditary hemolytic anemia	282.0; 282.1; 282.2; 282.3	D58
Dermatopolymyositis	716	M33
Myositis	717.90	M60
Dermatitis herpetiformis	693.00	L13.0
Henoch schoenlein purpura	287.00	D69.0
Idiopathic fibrosing alveolitis	517.01	J84
Idiopathic thrombocytopenic purpura	287.10	D69.3
Iridocyclitis	364.00	H20
Juvenile arthritis (nonsystemic)	712.00	M08.1, M08.3–8
LED (lupus erythematosus discoids)	695.40	L93.0
Mixed connective tissue disease (MCTD)	734.91	M35.1; M35.8; M35.9
Myasthenia gravis	733.00	G70.0
Pernicious anemia	281.00; 281.09	D51.0
Primary adrenocortical insufficiency, other adrenal gland	255.10	E27.1
Pemphigus	694.00–04, 694.08–09	L10
Pemphigoid	—	L12
Polyarteritis nodosa (microscopic polyangiitis)	446.00	M30
Churg strauss syndrome (eosinophilic granulomatosis with polyangiitis)	446.40	M30.1
Granulomatosis with polyangiitis (Wegener's granulomatosis/vasculitides)	446.29–0.31; 446.39	M31.3, M31.5, M31.6, M35.3
Seropositive rheumatoid arthritis	712.19, 712.39, 712.59	M05; M06
Other rheumatoid arthritis	712.30	M06
Psoriasis, vulgaris, psoriatic arthritis, generalized familial pustular psoriasis	696.09; 696.10, 696.19, 696.0–0.1	L40.0–3, L40.5–9
Sarcoidosis	135.99	D86
Scleroderma	734	M34
Seropositive rheumatoid arthritis	712.19, 712.39, 712.59	M05; M06
Other rheumatoid arthritis	712.30	—
Uveitis	366.00	H20
Vitiligo	709.01	L80
Rheumatic fever/rheumatic myocarditis	390/391	I01
Glomerulonephritis	580–583	N00–N01.9, N03, N05
Kawasaki disease	—	M30.3
Raynaud's syndrome	—	I73.0
Interstitial cystitis	595.00	N30.1
Endometriosis	625.30	N80
Autoimmune hepatitis	571.93	K73
Autoimmune thyroiditis	245.03	E06.3
Autoimmune encephalitis	—	G04, G05.8
Celiac disease	269.00	K90.0
Guillain-Barre syndrome	354.00	G61.0
Multiple sclerosis	340.00	G35
Sjögren's syndrome	734.90	M35.0
Systemic lupus erythematosus	734.10	M32.1, M32.9
Thyrotoxicosis (Graves' disease)	242.00	E05.0
Type 1 diabetes	250.00	E10
Gastrointestinal involvement		
Regional enteritis (Crohn's disease)	563.00–0.02; 563.08; 563.09	K50
Autoimmune hepatitis, chronic active hepatitis	571.93	K73
Ulcerative colitis	563.10	K51
Primary biliary cirrhosis	571.90	K74.3
Celiac disease	269.00	K90.0
Autoinflammatory diseases		

TABLE 5 Continued

Autoimmune Diseases	ICD-8	ICD-10
Familial cold-autoinflammatory syndrome 2 (FCAS2-NLRP12)	—	L50.2
Neonatal-onset multisystem inflammatory disease (NOMID)	—	G03.1; G44.8; M08.9
Cherubism (SH3BP2)	526.90	K10.8 c + d
Behçet syndrome	—	M35.2
Recurrent pericarditis	—	I09.2
Chronic recurrent multifocal osteomyelitis (CRMO)	720.10	M86.3
Schnitzler syndrome, Muckle-Wells syndrome (MWS)	708.80	L50.8
Age-related macular degeneration	—	H35.3
Hemolytic uremic syndrome	—	D59.3

WHO, World Health Organization; —, not applicable.