

Supplemental Information

SUPPLEMENTAL TABLE 6 ICD-9 Codes of Diagnostic Groups and Self-Harm Methods

Diagnostic Group	ICD-9 Codes
ADHD	314.00, 314.01, 314.9
Disruptive behavior disorders	312.81, 312.9, 313.81, 312.34
Depressive disorders	296.2×, 296.3×, 296.82, 298.0, 300.4, 311
Bipolar disorders	296.0×, 296.1×, 296.4×, 296.5×, 296.6×, 296.7×, 296.8× (other than 296.82)
Anxiety disorders	300.0×, 300.2×, 300.3 308.3, 309.81, 293.84, 300.00– 300.02, 300.09, 300.20–300.23, 300.29, 300.3, 300.7, 308.0–308.9, 309.21, 309.81, 312.39, 313.0, 313.21, 313.23
Schizophrenia and related psychoses	295.00–295.99 298.9
Substance use disorders	291, 292, 303, 304, 305
Personality disorders	301
Other mental disorders	290–319 not included in other diagnostic groups
Deliberate self-harm method	
Violent	
Firearm	E955.0–E955.5, E955.9
Other violent	E954; E953.0–E953.9; E957.0–E957.9; E958.1, E958.2, E958.3, E958.5, E958.6, E958.7
Nonviolent	
Poisoning	E950.0–E952.9
Cutting	E956
Other	E955.6, E955.7, E958.0, E958.4, E958.8, E958.9

Diagnostic groups define diagnostic categories. At least 1 inpatient or at least 2 outpatient codes with a diagnostic group was required during 180 days before self-harm event for diagnostic groups. ICD-9, *International Classification of Diseases, Ninth Revision*.