



Supplemental Information



SUPPLEMENTAL FIGURE 2

Accrual table.



SUPPLEMENTAL FIGURE 3

The distribution of past-30-day cigarette use at follow-up by demographics (ie, sex, race, and highest parental education) and e-cigarette use in the overall sample. The points (jittered for clarity) show the number of past-30-day cigarettes smoked at follow-up by demographics along with violin plots and boxplots describing corresponding distributions. A, Overall sample. B, Past-30-day e-cigarette users at baseline.

SUPPLEMENTAL TABLE 3 The Frequencies and Percentages of Participants Who Were Included in the Analysis and Those Who Were Lost to Follow-up

	Analytic Sample, <i>N</i> (%)	Lost to Follow-up, <i>N</i> (%)	<i>P</i> for Difference ^a
<i>N</i>	1312	147	—
Sex			
Female	685 (52.2)	62 (42.2)	.026
Male	627 (47.8)	85 (57.8)	—
Race and/or ethnicity			
Non-Hispanic white	635 (48.4)	78 (53.1)	.520
Hispanic white	506 (38.6)	50 (34.0)	—
Other	171 (13.0)	19 (12.9)	—
Highest parental education			
≤12th grade	391 (29.8)	40 (27.2)	.007
Some college	459 (35.0)	60 (40.8)	—
College degree or higher	386 (29.4)	30 (20.4)	—
Missing	76 (5.8)	17 (11.6)	—
E-cigarette use			
Never	828 (63.1)	97 (66.0)	.036
Previous	346 (26.4)	27 (18.4)	—
Past 30 d	138 (10.5)	23 (15.6)	—

—, not applicable.

^a χ^2 tests of independence were conducted to calculate the *P* values.