

Supplemental Information

SUPPLEMENTAL TABLE 3 Results of Logistic Regression Modeling

	B	SE	P	Odds Ratio (95% CI)	Model Evaluation
Depressive disorders					$\chi^2_{LR} (5) = 20.48, P = .001$ $\chi^2_{HL} (8) = 3.50, P = .90$
Age of first visit	0.176	0.104	.09	1.20 (0.97–1.46)	
Date cohort 1	0.187	0.272	.47	1.21 (0.71–2.05)	
Pubertal stage late	1.703	0.800	.03	5.49 (1.14–26.32)	
Socially transitioned	0.167	0.329	.61	1.18 (0.62–2.25)	
Assigned sex female	0.237	0.367	.52	1.27 (0.62–2.60)	
Anxiety disorders ^a					$\chi^2_{LR} (5) = 13.42, P = .02$ $\chi^2_{HL} (8) = 5.16, P = .74$
Age of first visit	0.039	0.100	.70	1.04 (0.86–1.27)	
Date cohort 1	−0.310	0.264	.24	0.73 (0.44–1.23)	
Pubertal stage late	1.432	0.631	.02	4.18 (1.22–14.49)	
Socially transitioned	−0.034	0.315	.92	0.97 (0.52–1.80)	
Assigned sex female	0.283	0.353	.42	1.33 (0.67–2.65)	
History of self-harm					$\chi^2_{LR} (5) = 22.75, P < .001$ $\chi^2_{HL} (8) = 6.82, P = .56$
Age of first visit	0.185	0.109	.09	1.20 (0.97–1.49)	
Date cohort 1	0.006	0.280	.98	1.01 (0.58–1.74)	
Pubertal stage late	0.636	0.711	.37	1.89 (0.47–7.63)	
Socially transitioned	0.349	0.350	.32	1.42 (0.71–2.82)	
Assigned sex female	1.227	0.447	.006	3.41 (1.42–8.19)	
History of suicidal ideation					$\chi^2_{LR} (5) = 9.97, P = .08$ $\chi^2_{HL} (8) = 5.25, P = .73$
Age of first visit	0.190	0.099	.05	1.21 (1.00–1.47)	
Date cohort 1	−0.135	0.263	.61	0.87 (0.52–1.46)	
Pubertal stage late	0.478	0.543	.38	1.61 (0.56–4.67)	
Socially transitioned	0.127	0.310	.68	1.14 (0.62–2.08)	
Assigned sex female	0.115	0.342	.74	1.12 (0.57–2.19)	
Use of psychoactive medications					$\chi^2_{LR} (5) = 20.83, P = .001$ $\chi^2_{HL} (8) = 2.24, P = .97$
Age of first visit	0.269	0.113	.02	1.31 (1.05–1.63)	
Date cohort 1	0.109	0.287	.70	1.12 (0.64–1.96)	
Pubertal stage late	0.909	0.815	.26	2.48 (0.50–12.20)	
Socially transitioned	0.298	0.359	.41	1.35 (0.67–2.72)	
Assigned sex female	0.690	0.418	.10	2.00 (0.88–4.53)	

The above models met conditions for goodness-of-fit testing. Models for current suicidal ideation and previous suicide attempts are not included because conditions for goodness-of-fit testing were not met. χ^2_{HL} , Hosmer-Lemeshow goodness-of-fit test statistic; χ^2_{LR} , likelihood ratio test statistic; —, not applicable.

^a Generalized anxiety disorder, panic disorder, social anxiety, and obsessive-compulsive disorder are included.