

## Supplemental Information

**SUPPLEMENTAL TABLE 3** Sensitivity Analysis of Prevalence of STI Testing (With 'Unsure' Respondents Categorized as 'Yes') Other Than HIV Testing in the Past 12 Months by Demographic Characteristics Among Currently Sexually Active US High School Students by Sex, 2019

	Prevalence of STI Testing in Past 12 Mo		
	Total, % (95% CI) (N = 2501) <sup>a</sup>	Males, % (95% CI) (n = 1178) <sup>a</sup>	Females, % (95% CI) (n = 1301) <sup>a</sup>
Total <sup>b</sup>	24.5 (21.3–28.0)	18.8 (15.8–22.2)	29.4 (25.2–34.0)
Race and ethnicity <sup>c</sup>			
Non-Hispanic White	22.9 (19.2–26.9)	14.0 (10.5–18.6)	30.0 (24.6–36.0)
Non-Hispanic Black	29.3 (20.4–40.0)	22.2 (13.4–34.6)	38.6 (26.5–52.2)
Hispanic	24.0 (18.3–30.8)	21.9 (16.3–28.8)	25.8 (18.8–34.4)
Other <sup>d</sup>	28.1 (19.9–38.1)	27.1 (17.9–38.8)	28.2 (17.3–42.3)
Age, y <sup>e</sup>			
≤15	20.4 (15.3–26.6)	24.4 (17.0–33.8)	16.0 (10.5–23.6)
16	21.0 (16.1–27.0)	15.4 (10.4–22.3)	25.9 (20.0–32.9)
17	26.3 (21.4–31.8)	20.0 (15.7–25.3)	31.2 (24.2–39.2)
≥18	28.8 (24.5–33.5)	16.9 (12.2–22.9)	41.1 (33.6–49.0)
Sexual identity			
Heterosexual	23.0 (19.9–26.4)	17.7 (15.1–20.7)	28.6 (23.6–34.3)
Lesbian, gay, bisexual	32.2 (25.8–39.4)	30.8 (14.0–55.0)	32.7 (26.5–39.6)
Unsure	36.7 (24.2–51.3)	—	27.7 (14.9–45.6)
Sex of sexual contact			
Same sex or both sex	34.9 (27.5–43.3)	39.4 (21.7–60.4)	33.7 (26.6–41.7)
Opposite sex only	22.8 (19.8–26.2)	17.6 (14.7–20.8)	28.3 (23.8–33.2)
Had sexual intercourse before age 13 <sup>d,e</sup>			
Yes	39.2 (29.2–50.3)	34.5 (24.6–46.1)	48.5 (33.0–64.3)
No	23.4 (20.4–26.7)	16.9 (14.1–20.1)	28.8 (24.6–33.3)
Had sexual intercourse with 2 or more persons during the previous 3 mo <sup>d</sup>			
Yes	31.6 (25.8–38.0)	28.6 (22.1–36.1)	35.4 (27.1–44.6)
No	22.6 (19.6–26.0)	15.0 (12.6–17.9)	28.4 (24.1–33.0)
Had ≥4 lifetime sexual partners <sup>d</sup>			
Yes	32.4 (27.2–38.1)	29.2 (23.9–35.0)	35.8 (27.8–44.8)
No	21.7 (18.6–25.1)	13.7 (10.9–16.9)	27.8 (23.4–32.6)
Did not use a condom at last sexual intercourse <sup>d</sup>			
Yes	30.7 (25.9–36.0)	19.6 (15.5–24.4)	38.6 (32.1–45.5)
No	19.7 (16.5–23.2)	18.8 (14.8–23.6)	20.2 (15.9–25.2)
Used alcohol or drugs before last sexual intercourse <sup>d,e</sup>			
Yes	30.1 (24.2–36.8)	24.8 (17.8–33.4)	36.6 (29.0–45.0)
No	22.7 (19.6–26.2)	16.8 (14.1–19.9)	27.5 (23.1–32.4)

Currently, sexually active is defined as those who had sexual intercourse with at least 1 person during the 3 mo before survey administration. STI, sexually transmitted infection; —, prevalence estimate suppressed due to insufficient sample size.

<sup>a</sup> Sample *n* is unweighted; percentage is weighted.

<sup>b</sup> Differences between males and females at *P* < .05.

<sup>c</sup> Other race category includes American Indian/Alaska Native, Asian American, Native Hawaiian/other Pacific Islander, and multiple non-Hispanic.

<sup>d</sup>  $\chi^2$  test statistic used to identify differences by STI testing status had a *P* value < .05 for males.

<sup>e</sup>  $\chi^2$  test statistic used to identify differences by STI testing status had a *P* value < .05 for females.