

Supplemental Information

SUPPLEMENTAL TABLE 2 Images Obtained and Changes in Methemoglobin and Hemoglobin for Each Patient

Case	Imaging Tests Performed	Methemoglobin			Hemoglobin		
		Before Dapsone Treatment (%)	During Treatment: Peak (%)	Time to Peak, d	Before Dapsone Treatment (g/dL)	During Treatment: Nadir (g/dL)	Time to Nadir, d
1	US	0.7	3.2	44	13.5	10.3	65
2	US, CT, EGD	0.3	3.5	14	9.6	9.2	7
3	US	0.6	6.4	70	13.7	10.5	62
4	US	0.3	4.2	5	15.1	11.7	79
5	US, CT	0.4	5.0	6	13.7	11.0	135
6	US, EGD	0.2	3.1	5	14.6	12.5	69
7	US, EGD, SBCE	0.7	5.0	21	13.6	9.4	21
Median (range)		0.4 (0.2 to 0.7)	4.2 (3.1 to 6.4)	14 (5 to 70)	13.7 (9.6 to 15.1)	10.5 (9.2 to 12.5)	65 (7 to 135)

US, ultrasound; CT, computed tomography; EGD, esophago-gastroduodenoscopy; SBCE, small-bowel capsule endoscopy; —, not applicable.